

## On Generalize Some Weak Forms of Supra Mappings in Intuitionistic Topological Spaces

Samer, R. Yaseen<sup>1</sup>, Hind, F. Abbas

<sup>1</sup> Department of Mathematics, College of Science, Tikrit University, Tikrit, Iraq

<sup>2</sup> Khaled Ibn Al Walid School, Directorate of Education Salah Eddin, Tikrit, Iraq

### ARTICLE INFO.

#### Article history:

-Received: 6 / 12 / 2017

-Accepted: 21 / 1 / 2018

-Available online: / / 2018

**Keywords:** Intuitionistic Topology, Intuitionistic, Generalized Pre supra mapping, Intuitionistic Generalized Semi supra mapping, Intuitionistic Generalized  $\alpha$ -supra mapping, Intuitionistic Generalized  $\beta$ -supra mapping.

#### Corresponding Author:

Name: Samer, R. Yaseen

E-mail: [amer2017@tu.edu.iq](mailto:amer2017@tu.edu.iq)

Tel:

### 1. Introduction

Zadeh [1] was introducing the concept of "fuzzy set" in 1965. After that in 1968 Chang [2] introduced the concept of "fuzzy topology". Also in 1983, Atanassov introduced the concept of "Intuitionistic fuzzy set" [3,4]. Finally Coker [5], introduced concept of "intuitionistic sets" and using it to introduce the concept of "intuitionistic topological spaces" [6].

In this paper we generalize some weak forms of supra mappings in Intuitionistic topological spaces and studied some of their properties and relationships among them.

### 2. Preliminaries

**Definition 2.1** [5] Let  $M \subseteq X \neq \emptyset$  and. The Intuitionistic set  $\tilde{M}$  (IS, for short) is the form  $\tilde{M} = \langle x, M_1, M_2 \rangle$  and  $M_1, M_2 \subseteq X$  with condition  $M_1 \cap M_2 = \emptyset$ . The set  $M_1$  is called "the set of members" of  $\tilde{M}$  and  $M_2$  is called "the set of non-members" of  $\tilde{M}$ .

### Abstract

The aim of this paper is to introduce a new classes of supra mappings called Intuitionistic Generalized Pre supra mapping, Intuitionistic Generalized Semi supra mapping, Intuitionistic Generalized  $\alpha$ - supra mapping and Intuitionistic Generalized  $\beta$ -supra mapping. At last we studied some of their properties and investigate relationships among this concepts.

**Definition 2. 2** [5] Let  $X \neq \emptyset$ , and let  $\tilde{M} = \langle x, M_1, M_2 \rangle$ ,  $\tilde{N} = \langle x, N_1, N_2 \rangle$  are two Intuitionistic sets respectively. Also, let  $\{\tilde{M}_s; s \in S\}$  be a collection of "Intuitionistic sets in  $X$ ", and  $\tilde{M}_i = \langle x, M_s^{(1)}, M_s^{(2)} \rangle$ , the following is valid.

- 1)  $\tilde{M} \tilde{M} \subseteq \tilde{N}$  iff  $M_1 \subseteq N_1$  and  $N_2 \subseteq M_2$ ,
- 2)  $\tilde{M} = \tilde{N}$  iff  $\tilde{M} \subseteq \tilde{N}$  and  $\tilde{N} \subseteq \tilde{M}$ ,
- 3) The complement of  $\tilde{M}$  is denoted by  $\bar{\tilde{M}}$  and defined by  $\bar{\tilde{M}} = \langle x, M_2, M_1 \rangle$ ,
- 4)  $\cup \tilde{M}_i = \langle x, \cup M_s^{(1)}, \cap M_s^{(2)} \rangle$ ,  $\cap \tilde{M}_i = \langle x, \cap M_s^{(1)}, \cup M_s^{(2)} \rangle$ ,
- 5)  $\emptyset = \langle x, \emptyset, X \rangle$ ,  $\bar{X} = \langle x, X, \emptyset \rangle$ .

**Definition 2.3** [6] Let  $X \neq \emptyset$ ,  $w \in X$  and let  $\tilde{M} = \langle x, M_1, M_2 \rangle$  be an Intuitionistic set the Intuitionistic point (IP f, for briefly) "Is  $w$ " defined by  $\dot{w} = \langle x, \{w\}, \{w\}^c \rangle$  in  $X$ . Also a Vanishing Intuitionistic point defined by  $Is \dot{w} = \langle x, \emptyset, \{w\}^c \rangle$  in  $X$ . The  $Is \dot{w}$  is said belong in  $\tilde{M}$  ( $\dot{w} \in M$ , for brief) iff  $w \in M_1$ , also  $Is \ddot{w}$  contained in  $\tilde{M}$

$(\tilde{w} \in \tilde{M}, \text{ for short})$  iff  $w \notin M_2$ .

**Definition 2.4 [5]** Let  $X, Y \neq \emptyset$ , and  $r : (X, \mu) \rightarrow (Y, \gamma)$  be a mapping.

a) If  $\tilde{N} = \langle y, N_1, N_2 \rangle$  is an Is in  $Y$ , then the inverse image of  $\tilde{N}$  under  $r$  defined by

$$r^{-1}(\tilde{N}) = \langle x, r^{-1}(N_1), r^{-1}(N_2) \rangle.$$

b) If  $\tilde{M} = \langle x, M_1, M_2 \rangle$  is an Is in  $X$ , then  $r(\tilde{V}) = \langle y, r(M_1), \tilde{r}(M_2) \rangle$  is an Is in  $Y$

$$\text{where } \tilde{r}(\tilde{M}) = \overline{r(\tilde{M}_2)}.$$

**Definition 2.5 [7]** Let  $X \neq \emptyset$ . An Intuitionistic topology (ITS, for short) on  $X$  is

a collection  $\mu$  of an "Intuitionistic sets" in  $X$  satisfying :

- (1)  $\emptyset, X \in \mu$ .
- (2)  $\mu$  is closed under finite intersections.
- (3)  $\mu$  is closed under arbitrary unions.

Each element in  $\mu$  is called "Intuitionistic open set" and denoted by "IOS"

The complement of an "Intuitionistic open set" is called "Intuitionistic closed set" denoted by "ICS".

**Definition 2.6 [7]** Let  $(X, \mu)$  be an ITS and let  $\tilde{M} = \langle x, M_1, M_2 \rangle \subseteq X$ . The

"interior" (namely,  $\text{int}(\tilde{M})$ ) and the "closure" (namely,  $\text{cl}(\tilde{M})$ ) are defined :

$$\text{int}(\tilde{M}) = \cup \{ \tilde{V} : \tilde{V} \subseteq \tilde{M}, \tilde{V} \in \mu \},$$

$$\text{cl}(\tilde{M}) = \cap \{ \tilde{J} : \tilde{M} \subseteq \tilde{J}, \tilde{J} \in \mu \}. \text{ Also}$$

$$1\text{-sint}(\tilde{M}) = \cup \{ \tilde{V} : \tilde{V} \subseteq \tilde{M}, \tilde{V} \in \text{ISOX} \},$$

$$\text{scl}(\tilde{M}) = \cap \{ \tilde{J} : \tilde{M} \subseteq \tilde{J}, \tilde{J} \in \text{ISCS} \}.$$

$$2\text{-pint}(\tilde{M}) = \cup \{ \tilde{V} : \tilde{V} \subseteq \tilde{M}, \tilde{V} \in \text{IPOX} \},$$

$$\text{pcl}(\tilde{M}) = \cap \{ \tilde{J} : \tilde{M} \subseteq \tilde{J}, \tilde{J} \in \text{IPCS} \}.$$

$$3\text{-}\alpha\text{int}(\tilde{M}) = \cup \{ \tilde{V} : \tilde{V} \subseteq \tilde{M}, \tilde{V} \in \text{I}\alpha\text{OX} \},$$

$$\alpha\text{cl}(\tilde{M}) = \cap \{ \tilde{J} : \tilde{M} \subseteq \tilde{J}, \tilde{J} \in \text{I}\alpha\text{CS} \}.$$

$$4\text{-}\beta\text{int}(\tilde{M}) = \cup \{ \tilde{V} : \tilde{V} \subseteq \tilde{M}, \tilde{V} \in \text{I}\beta\text{OX} \},$$

$$\beta\text{cl}(\tilde{M}) = \cap \{ \tilde{J} : \tilde{M} \subseteq \tilde{J}, \tilde{J} \in \text{I}\beta\text{CS} \}.$$

**Remark 2-7.[7]** This implications are valid :

$\text{sint}(\tilde{M}) \subseteq \tilde{M}$ ,  $\text{scl}(\tilde{M}) = \tilde{M}$ ,  $\text{pint}(\tilde{M}) \subseteq \tilde{M}$ ,  $\text{pcl}(\tilde{M}) = \tilde{M}$ ,  $\alpha\text{int}(\tilde{M}) \subseteq \tilde{M}$ ,  $\alpha\text{cl}(\tilde{M}) = \tilde{M}$ ,  $\beta\text{int}(\tilde{M}) \subseteq \tilde{M}$  and  $\beta\text{cl}(\tilde{M}) = \tilde{M}$ .

**Definition 2.8. [8]**

Let  $(X, \mu)$  be an ITS. IS  $\tilde{M}$  of  $X$  is said to be

1. ISOS if  $\tilde{M} \subseteq \text{Icl}(\text{Iint}(\tilde{M}))$ ,
2. IPOS if  $\tilde{M} \subseteq \text{Iint}(\text{Icl}(\tilde{M}))$ ,
3. I $\alpha$ OS if  $\tilde{M} \subseteq \square \text{Iint}(\text{Icl}(\text{Iint}(\tilde{M})))$ ,
4. I $\beta$ OS if  $\tilde{M} \subseteq \text{Icl}(\text{Iint}(\text{Icl}(\tilde{M})))$ .

The family of all intuitionistic semi-open, pre-open,  $\alpha$ -open and  $\beta$ -open sets of  $(X, \mu)$  are denoted by "ISOS(X)", "IPOS(X)", "I $\alpha$ OS(X)" and "I $\beta$ OS(X)" respectively. Also the complement of all intuitionistic semi-open, pre-open,  $\alpha$ -open and  $\beta$ -open sets of  $(X, \mu)$  are denoted by "ISCS(X)", "IPCS(X)", "I $\alpha$ CS(X)" and "I $\beta$ CS(X)" respectively.

**Definition 2.9. [8]**

Let  $(X, \mu)$  be an ITS. An intuitionistic set  $\tilde{M}$  of  $X$  is said to be :

1) Intuitionistic generalizes-open set (Igos, for short) if  $\forall U$  is ICS s.t  $U \subseteq \tilde{M}$  then  $U \subseteq \text{int}(\tilde{M})$ .

2) Intuitionistic generalizes semi-open set (Igsos, for short) if  $\forall U$  is ISCS s.t  $U \subseteq \tilde{M}$  then  $U \subseteq \text{int}(\tilde{M})$ .

3) Intuitionistic generalizes pre-open set (IgpOs, for short) if  $\forall U$  is IPCS s.t  $U \subseteq \tilde{M}$  then  $U \subseteq \text{int}(\tilde{M})$ .

4) Intuitionistic generalizes  $\alpha$ -open set (I $\alpha$ Os, for short) if  $\forall U$  is I $\alpha$ CS s.t  $U \subseteq \tilde{M}$  then  $U \subseteq \text{int}(\tilde{M})$ .

5) Intuitionistic generalizes  $\beta$ -open set (I $\beta$ Os, for short) if  $\forall U$  is I $\beta$ CS s.t  $U \subseteq \tilde{M}$  then  $U \subseteq \text{int}(\tilde{M})$ .

**Definition 2.10. [8]**

A map  $r : (X, \mu) \rightarrow (Y, \delta)$  is said to be:

1- intuitionistic continuous if the pre-image  $f^{-1}(\tilde{M})$  is IOS in  $X$  for every IOS  $\tilde{M}$  in  $Y$ .

2. Intuitionistic pre continuous if the pre image  $f^{-1}(\tilde{M})$  is IPOS in  $X$  for every IOS  $\tilde{M}$  in  $Y$ .

3. Intuitionistic semi continuous if the pre image  $f^{-1}(\tilde{M})$  is ISOS in  $X$  for every IOS  $\tilde{M}$  in  $Y$ .

4- Intuitionistic  $\alpha$ -continuous if the pre image  $f^{-1}(\tilde{M})$  is I $\alpha$ OS in  $X$  for every IOS  $\tilde{M}$  in  $Y$ .

5- Intuitionistic  $\beta$ -continuous if the pre image  $f^{-1}(\tilde{M})$  is I $\beta$ OS in  $X$  for every IOS  $\tilde{M}$  in  $Y$ .

Now, we give this definition.

**Definition 2.11.**

A map  $r : (X, \mu) \rightarrow (Y, \delta)$  is said to be:

1. IgP continuous mapping if the pre image  $f^{-1}(\tilde{M})$  is IPCS in  $X$  for every IOS  $\tilde{M}$  in  $Y$ .

2. IgS continuous mapping if the pre image  $f^{-1}(\tilde{M})$  is ISCS in  $X$  for every IOS  $\tilde{M}$  in  $Y$ .

3- Ig $\alpha$ -continuous mapping if the pre image  $f^{-1}(\tilde{M})$  is I $\alpha$ CS in  $X$  for every IOS  $\tilde{M}$  in  $Y$ .

4- Ig $\beta$ -continuous mapping if the pre image  $f^{-1}(\tilde{M})$  is I $\beta$ CS in  $X$  for every IOS  $\tilde{M}$  in  $Y$ .

## Section 2 INTUITIONISTIC GENERALIZED PRE, SEMI, $\beta$ & $\alpha$ - of SUPRA MAPPINGS

In this section we have introduced intuitionistic this concepts: generalized pre supra mapping, Intuitionistic generalized semi supra mapping, Intuitionistic generalized  $\beta$  supra mappings, Intuitionistic generalized  $\alpha$ - supra mapping and studied some from its properties.

**Definition 2.1:** A mapping  $r : (X, \mu) \rightarrow (Y, \gamma)$  be an "Intuitionistic generalized pre supra mapping" ("IgpSm", for short) (resp., "Intuitionistic generalized semi supra mapping" ("IgsSm", for short), "Intuitionistic generalized  $\alpha$  supra mapping" (I $\alpha$ asm, for short), "Intuitionistic generalized  $\beta$  supra mapping" (I $\beta$ asm, for short) if  $r^{-1}(\tilde{M})$  is an "IgpOS" (resp., is an "IgsOS", "I $\alpha$ OS", "I $\beta$ OS")

in  $(X, \mu)$  for every "IgPOS"  $\tilde{M}$  of  $(Y, \gamma)$  ((resp., for every "IgSOS"  $\tilde{M}$ , "Ig $\alpha$ OS"  $\tilde{M}$  "Ig $\beta$ OS"  $\tilde{M}$  of  $(Y, \gamma)$ ).

**Proposition 2.2:** Let  $r: (E, \mu) \rightarrow (D, \gamma)$  and  $p: (D, \gamma) \rightarrow (J, \delta)$  be IgPsm. Then  $p \circ r: (E, \mu) \rightarrow (J, \delta)$  is IgPsm.

**Proof:** Let  $\tilde{M}$  be IgPOS in  $J$ . Then  $p^{-1}(\tilde{M})$  is IgPOS in  $D$ , since  $r$  is IgPsm, so  $r^{-1}(p^{-1}(\tilde{M}))$  is IgPOS in  $E$ . Therefore  $p \circ r$  is an IgPsm.

**Proposition 2.3:** Let  $r: (E, \mu) \rightarrow (D, \gamma)$  and  $p: (D, \gamma) \rightarrow (J, \delta)$  be Ig $\alpha$ sm. Then  $p \circ r: (E, \mu) \rightarrow (J, \delta)$  is IgSsm.

**Proof:** Let  $\tilde{M}$  be Ig $\alpha$ OS in  $J$ . So that  $p^{-1}(\tilde{M})$  is Ig $\alpha$ OS in  $D$ , since  $r$  is Ig $\alpha$ sm and every Ig $\alpha$ OS is IgSOS. Thus  $r^{-1}(p^{-1}(\tilde{M}))$  is IgSOS in  $E$ . Therefore  $p \circ r$  is IgSsm.

**Proposition 2.4:** : Let  $r: (E, \mu) \rightarrow (D, \gamma)$  be an IgPsm and  $p: (D, \gamma) \rightarrow (J, \delta)$  be IgP continuous supra mapping, then  $p \circ r: (E, \mu) \rightarrow (J, \delta)$  is IgP continuous supra mapping.

**Proof:** Let  $\tilde{M}$  be IOS in  $J$ . Then  $p^{-1}(\tilde{M})$  is IgPOS in  $Y$ . Since  $r$  is IgPsm, then  $r^{-1}(p^{-1}(\tilde{M}))$  is IgPOS in  $X$ . Therefore  $p \circ r$  is IgP continuous supra mapping.

**Proposition 2.5:** : Let  $r: (E, \mu) \rightarrow (D, \gamma)$  be an Igsm and  $p: (D, \gamma) \rightarrow (J, \delta)$

be IgP continuous supra mapping, then  $p \circ r: (E, \mu) \rightarrow (J, \delta)$  is IgP continuous supra mapping.

**Proof:** it's obvious.

**Proposition 2.6:** Let  $r: (E, \mu) \rightarrow (D, \gamma)$  be an Ig $\alpha$ sm and  $p: (D, \gamma) \rightarrow (J, \delta)$

be Ig $\alpha$  continuous mapping, then  $p \circ r: (E, \mu) \rightarrow (J, \delta)$  is an IgS continuous mapping.

**Proof:** Let  $\tilde{M}$  be IOS in  $J$ . Then  $p^{-1}(\tilde{M})$  is an Ig $\alpha$ OS in  $D$ . Since  $r$  is Ig $\alpha$ sm, then  $r^{-1}(p^{-1}(\tilde{M}))$  is IgSOS in  $E$  and every Ig $\alpha$ OS is IgSOS. Hence  $p \circ r$  is IgS continuous mapping.

**Proposition 2.7:** Let  $r: (E, \mu) \rightarrow (D, \gamma)$  be IgPsm and  $p: (D, \gamma) \rightarrow (J, \delta)$

be IgP continuous mapping, then  $p \circ r: (E, \mu) \rightarrow (J, \delta)$  is Ig $\beta$  continuous mapping.

**Proof:** it's obvious.

**Proposition 2.8:** Let  $r: (E, \mu) \rightarrow (D, \delta)$  be Ig $\alpha$ sm. Then this implications are equivalent:

- (i)  $r^{-1}(\tilde{M})$  is IgSOS in  $E$  for each IgSOS  $\tilde{M}$  in  $D$ ,
- (ii)  $r^{-1} \alpha \text{int}(\tilde{M}) \subseteq \text{sint } r^{-1}(\tilde{M})$  for every "Is  $\tilde{M}$ " of  $D$ ,
- (iii)  $\alpha \text{cl } r^{-1}(\tilde{M}) \subseteq r^{-1} \text{scl}(\tilde{M}) \forall$  "Is  $\tilde{M}$ " of  $D$ .

**Proof:** (i)  $\Rightarrow$  (ii) Let  $\tilde{M}$  be ISOS in  $D$  and  $\alpha \text{int}(\tilde{M}) \subseteq \tilde{M}$ , so  $r^{-1} \alpha \text{int}(\tilde{M}) \subseteq r^{-1}(\tilde{M})$ . Since  $\alpha \text{int}(\tilde{M})$  is I $\alpha$ OS in  $D$ , and every I $\alpha$ OS is ISOS. So IgSOS in  $D$ . Therefore  $r^{-1} \alpha \text{int}(\tilde{M})$  is IgSOS in  $E$ , and  $r^{-1} \alpha \text{int}(\tilde{M})$  is ISOS in  $E$ , since  $r^{-1} \alpha \text{int}(\tilde{M}) \subseteq r^{-1} \text{sint}(\tilde{M})$ .

Thus  $r^{-1} \alpha \text{int}(\tilde{M}) = \alpha \text{int } r^{-1} \alpha \text{int}(\tilde{M}) \subseteq \alpha \text{int } r^{-1}(\tilde{M}) \subseteq \text{sint } r^{-1} \text{sint}(\tilde{M}) \subseteq \text{sint } r^{-1}(\tilde{M})$ .

(ii)  $\Rightarrow$  (iii) by taking complement of (ii) we get the result of (iii).

(iii)  $\Rightarrow$  (i) Let  $\tilde{M}$  be IgSCS in  $D$ . Since  $\tilde{M}$  is ISCS in  $D$  and  $\text{scl}(\tilde{M}) = \tilde{M}$ . So  $r^{-1}(\tilde{M}) = \alpha \text{cl } r^{-1}(\tilde{M}) \subseteq r^{-1} \text{scl}(\tilde{M})$ . Therefore  $r^{-1}(\tilde{M})$  is IgSCS in  $E$ .

**Proposition 2.9:** Let  $r: (E, \mu) \rightarrow (D, \delta)$  be Ig $\alpha$ sm. Then this implications are equivalent:

- (i)  $r^{-1}(\tilde{M})$  is Ig $\alpha$ OS in  $E \forall$  Ig $\alpha$ OS  $\tilde{M}$  in  $D$ ,
- (ii)  $r^{-1} \text{pint}(\tilde{M}) \subseteq \beta \text{int } r^{-1}(\tilde{M})$  for every Is  $\tilde{M}$  of  $D$ ,
- (iii)  $\text{pcl } r^{-1}(\tilde{M}) \subseteq r^{-1} \beta \text{cl}(\tilde{M}) \forall$  IS  $\tilde{M}$  of  $D$ .

**Proof:** (i)  $\Rightarrow$  (ii) Let  $\tilde{M}$  be I $\alpha$ OS in  $D$  and  $\alpha \text{int}(\tilde{M}) \subseteq \tilde{M}$ , and  $r^{-1} \alpha \text{int}(\tilde{M}) \subseteq r^{-1}(\tilde{M})$ . Since  $\alpha \text{int}(\tilde{M})$  is I $\beta$ OS in  $D$ , and every I $\alpha$ OS is I $\beta$ OS. So Ig $\beta$ OS in  $D$ . Therefore  $r^{-1} \alpha \text{int}(\tilde{M})$  is Ig $\beta$ OS in  $E$ , and  $r^{-1} \alpha \text{int}(\tilde{M})$  is I $\beta$ OS in  $E$ , since  $r^{-1} \alpha \text{int}(\tilde{M}) \subseteq r^{-1} \text{pint}(\tilde{M}) \subseteq r^{-1} \beta \text{int}(\tilde{M})$ . Thus  $r^{-1} \text{pint}(\tilde{M}) = \text{pint } r^{-1} \text{pint}(\tilde{M}) \subseteq \text{pint } r^{-1}(\tilde{M}) \subseteq \beta \text{int } r^{-1} \beta \text{int}(\tilde{M}) \subseteq \beta \text{int } r^{-1}(\tilde{M})$ .

(ii)  $\Rightarrow$  (iii) by taking complement of (ii) we get the result of (iii).

(iii)  $\Rightarrow$  (i) Let  $\tilde{M}$  be Ig $\alpha$ OS in  $D$ , since  $\tilde{M}$  is I $\alpha$ OS in  $D$  and  $\alpha \text{cl}(\tilde{M}) = \tilde{M}$ . Thus

$r^{-1}(\tilde{M}) = \alpha \text{cl } r^{-1}(\tilde{M}) \subseteq r^{-1} \alpha \text{cl}(\tilde{M})$ . Therefore  $r^{-1}(\tilde{M})$  is Ig $\alpha$ OS in  $E$ .

**Proposition 2.10:** Let  $r: (E, \mu) \rightarrow (D, \delta)$  be IgPsm. Then this implications are equivalent:

- (i)  $r^{-1}(\tilde{M})$  is IgPOS in  $E \forall$  IgPOS  $\tilde{M}$  in  $D$ ,
- (ii)  $r^{-1} \text{pint}(\tilde{M}) \subseteq \beta \text{int } r^{-1}(\tilde{M})$  for every Is  $\tilde{M}$  of  $D$ ,
- (iii)  $\text{pcl } r^{-1}(\tilde{M}) \subseteq r^{-1} \beta \text{cl}(\tilde{M}) \forall$  IS  $\tilde{M}$  of  $D$ .

**Proof:** it's obvious.

**Proposition 2.11:** Let  $r: (E, \mu) \rightarrow (D, \gamma)$  and  $p: (D, \gamma) \rightarrow (J, \delta)$  are two Ig $\alpha$ sm. Then

$p \circ r: (E, \mu) \rightarrow (J, \delta)$  is IgPsm.

**Proof:** Let  $\tilde{M}$  be Ig $\alpha$ OS in  $J$ . Thus  $p^{-1}(\tilde{M})$  is Ig $\alpha$ OS in  $D$ , since  $r$  is Ig $\alpha$ sm, then  $r^{-1}(p^{-1}(\tilde{M}))$  is Ig $\alpha$ OS in  $E$ . Since every Ig $\alpha$ OS is IgPOS. Thus  $r^{-1}(p^{-1}(\tilde{M}))$  is IgPOS in  $E$ . Therefore  $p \circ r$  is IgPsm.

**Proposition 2.12:** Let  $r: (E, \mu) \rightarrow (D, \gamma)$  be IgPsm and  $p: (D, \gamma) \rightarrow (J, \delta)$

be Ig $\alpha$  continuous mapping, then  $p \circ r: (E, \mu) \rightarrow (J, \gamma)$  is Ig $\beta$  continuous mapping.

**Proof:** Let  $\tilde{M}$  be Ig $\alpha$ OS in  $J$ . So that  $p^{-1}(\tilde{M})$  is Ig $\alpha$ OS in  $D$ , since every Ig $\alpha$ OS is IgPOS and  $r$  is IgPsm, then  $r^{-1}(p^{-1}(\tilde{M}))$  is IgPOS in  $E$ . Since every IgPOS is Ig $\beta$ OS. Thus  $r^{-1}(p^{-1}(\tilde{M}))$  is Ig $\beta$ OS in  $E$ . Thus  $p \circ r$  is Ig $\beta$  continuous mapping.

**Proposition 2.13:** Let  $r: (E, \mu) \rightarrow (D, \gamma)$  be Ig $\alpha$ sm and

$p: (D, \gamma) \rightarrow (J, \delta)$  be Ig $\alpha$  continuous mapping, then  $p \circ r: (E, \mu) \rightarrow (J, \gamma)$  is Ig $\beta$  continuous mapping.

**Proof:** Let  $\tilde{M}$  be I $\alpha$ OS in  $J$ . So that  $p^{-1}(\tilde{M})$  is Ig $\alpha$ OS in  $D$ , since  $r$  is Ig $\alpha$ sm,  $r^{-1}(p^{-1}(\tilde{M}))$  is Ig $\alpha$ OS in  $E$ . Since every Ig $\alpha$ OS is Ig $\beta$ OS. Thus  $r^{-1}(p^{-1}(\tilde{M}))$  is Ig $\beta$ OS in  $E$ . Therefore  $p \circ r$  is Ig $\beta$  continuous mapping.

**Proposition 2.14:** Let  $r : (E, \mu) \rightarrow (D, \gamma)$  be IgSsm and  $p : (D, \gamma) \rightarrow (J, \delta)$  be Ig continuous mapping , then  $p \circ r : (E, \mu) \rightarrow (J, \delta)$  is IgS continuous mapping .

**Proof:** Let  $\tilde{M}$  be IOS in J . So that  $p^{-1}(\tilde{M})$  is IgSOS in D, since r is IgSsm, then  $r^{-1}(p^{-1}(\tilde{M}))$  is IgOS in E . Since every IgOS is IgSOS . Thus  $r^{-1}(p^{-1}(\tilde{M}))$  is IgSOS in E . Therefore  $p \circ r$  is IgS continuous mapping.

**Proposition 2.15:** Let  $r : (E, \mu) \rightarrow (D, \gamma)$  be IgPsm and  $p : (D, \gamma) \rightarrow (J, \delta)$  be Igp continuous mapping , then  $p \circ r : (E, \mu) \rightarrow (J, \delta)$  is Igβ continuous mapping .

**Proof:** Let  $\tilde{M}$  be IPOS in J.Thus  $p^{-1}(\tilde{M})$  is IgPOS in D, since r is IgPsm, then  $r^{-1}(p^{-1}(\tilde{M}))$  is IgPOS in E . Since every IgPOS is IgβOS . Hence  $r^{-1}(p^{-1}(\tilde{M}))$  is IgβOS in E . Therefore  $p \circ r$  is Igβ continuous mapping.

**Proposition 2.16:** Let  $r : (E, \mu) \rightarrow (D, \gamma)$  be Igasm and  $p : (D, \gamma) \rightarrow (J, \delta)$  be Igs continuous mapping , then  $p \circ r : (E, \mu) \rightarrow (J, \delta)$  is Igβ continuous mapping .

**Proof:** it obvious .

**Proposition 2.17:** Let  $r : (E, \mu) \rightarrow (D, \gamma)$  be Igβsm and  $p : (D, \gamma) \rightarrow (J, \delta)$  be Igα continuous mapping , then  $p \circ r : (E, \mu) \rightarrow (J, \delta)$  is Igβ continuous mapping .

**Proof:** Let  $\tilde{M}$  be IαOS in J.Thus  $p^{-1}(\tilde{M})$  is IgβOS in D, since r is Igβsm, then  $r^{-1}(p^{-1}(\tilde{M}))$  is IgβOS in E . Since every IgαOS is IgβOS . Hence  $r^{-1}(p^{-1}(\tilde{M}))$  is IgβOS in E . Therefore  $p \circ r$  is Igβ continuous mapping.

**Proposition 2.18:** Let  $r : (E, \mu) \rightarrow (D, \gamma)$  be Igasm and  $p : (D, \gamma) \rightarrow (J, \delta)$  be Ig continuous mapping , then  $p \circ r : (E, \mu) \rightarrow (J, \delta)$  is IgP continuous mapping .

**Proof:** it obvious .


**Proposition 2.19:** Let  $r : (E, \mu) \rightarrow (D, \gamma)$  be Igasm and  $p : (D, \gamma) \rightarrow (J, \delta)$  be Ig continuous mapping , then  $p \circ r : (E, \mu) \rightarrow (J, \delta)$  is IgS continuous mapping .

**Proof:** it obvious .

**Section 3 The RELATIONS AMONG INTUITIONISTIC GENERALIZED PRE SUPRA MAPPING, INTUITIONISTIC GENERALIZED β – SUPRA MAPPING, INTUITIONISTIC GENERALIZED SEMI SUPRA MAPPING AND INTUITIONISTIC GENERALIZED α – SUPRA MAPPING .**

Now, We give this important theorem .

**Theorem 3.1.** The implication among some types of mappings are given by the following diagram.



Let  $r : (E, \mu) \rightarrow (D, \delta)$  be a mapping and  $\tilde{M}$  be IPOS in D, since r is IgPsm , then  $r^{-1}(\tilde{M})$  is IgPOS in E . Since each IPO(Y) is IβO(Y) .Hence  $r^{-1}(\tilde{M})$  is IgβOS in E for each  $\tilde{M}$  be IβOS in D. Therefore r is Igβsm .

**Igsm implies Igasm**

Let  $r : (E, \mu) \rightarrow (D, \delta)$  be a mapping and  $\tilde{M}$  be IOS in D. Since r is Igsm , then  $r^{-1}(\tilde{M})$  is IgOS in E . Since each IO(Y) is IαO(Y) .So that  $r^{-1}(\tilde{M})$  is IαOS in E  $\forall \tilde{M}$  be IαOS in Y. Therefore r is Igasm .

**Igasm implies IgSsm**

Let  $r : (E, \mu) \rightarrow (D, \delta)$  be a mapping and let  $\tilde{M}$  be IαOS in D. Since r is Igasm , then  $r^{-1}(\tilde{M})$  is IgαOS in E . Since each IαO(Y) is ISO(Y) .Thus  $r^{-1}(\tilde{M})$  is IgSOS in E  $\forall \tilde{M}$  be ISOS in Y. Therefore r is Igasm .

**IgSsm implies Igβsm**

Let  $r : (E, \mu) \rightarrow (D, \delta)$  be a mapping and let  $\tilde{M}$  be ISOS in D, since r is IgSsm , then  $r^{-1}(\tilde{M})$  is IgSOS in E . Since each ISO(Y) is IβO(Y) .Hence  $r^{-1}(\tilde{M})$  is IgβOS in E  $\forall \tilde{M}$  be an IβOS in D. Therefore r is Igβsm .

**Igasm implies IgPsm**

Let  $r : (E, \mu) \rightarrow (D, \delta)$  be a mapping and let  $\tilde{M}$  be ISOS in D, since r is IgSsm , then  $r^{-1}(\tilde{M})$  is IgSOS in E . Since each ISO(Y) is IβO(Y) .Hence  $r^{-1}(\tilde{M})$  is IgβOS in E  $\forall \tilde{M}$  be an IβOS in D. Therefore r is Igβsm .

**Remark 3.2** By transitivity we get this result:

- 1- Igasm implies Igβsm
- 2- Igsm implies IgPsm
- 3- Igsm implies IgSsm

The converse of the Theorem 3.1. is not true, the following examples are shown the cases.

**Example 3.3.** Let  $E = \{a, m, n\}$  with topology  $\mu = \{\tilde{E}, \emptyset, \tilde{S}, \tilde{R}, \tilde{K}, \tilde{U}\}$  , where  $\tilde{S} = \langle e, \{w\}, \{t, z\} \rangle$ ,  $\tilde{R} = \langle e, \{w, z\}, \emptyset \rangle$ ,  $\tilde{K} = \langle e, \{w\}, \emptyset \rangle$ ,  $\tilde{U} = \langle e, \{w\}, \{z\} \rangle$  &  $D = \{5, 6, 7\}$  with topology  $\delta = \{\tilde{Y}, \emptyset, \tilde{W}, \tilde{Q}\}$  , where  $\tilde{W} = \langle d, \{5\}, \{6, 7\} \rangle$ ,  $\tilde{Q} = \langle d, \{5\}, \emptyset \rangle$  . Let a mapping

$r : (E, \mu) \rightarrow (D, \delta)$  defined by  $r(\{a\}) = \{5\}$ ,  $r(\{m\}) = \{7\}$ ,  $r(\{n\}) = \{6\}$  . Then

1- r is Igβsm , because  $\forall \tilde{M}$  be IgβOS in D,  $r^{-1}(\tilde{M})$  is IgβOS in E. But r is not IgPsm, because  $r^{-1}(\{5, 7\}) = \{a, m\}$  is not IgPOS in E .

2- Also r is Igβsm , because  $\forall \tilde{M}$  be IgβOS in D,  $r^{-1}(\tilde{M})$  is IgβOS in E .But r is not IgSsm, because  $r^{-1}(\{6, 7\}) = \{m, n\}$  is not IgSOS in E .



**Example 3.4.** Let  $E = \{c, d, e\}$  with topology  $\mu = \{\dot{E}, \emptyset, \tilde{M}, \tilde{N}\}$ , where  $\tilde{M} = \langle e, \{c\}, \{d, e\} \rangle$ ,  $\tilde{N} = \langle e, \{c\}, \emptyset \rangle$  &  $D = \{1, 2, 3\}$  with topology  $\delta = \{\dot{D}, \emptyset, \tilde{O}, \tilde{F}\}$ , where  $\tilde{O} = \langle d, \{1\}, \{3\} \rangle$ ,  $\tilde{F} = \langle d, \{1\}, \emptyset \rangle$ . Let a mapping  $r : (E, \mu) \rightarrow (D, \delta)$  defined by  $r(\{c\}) = \{2\}$ ,  $r(\{d\}) = \{3\}$ ,  $r(\{e\}) = \{1\}$ . Thus  $r$  is IgPsm, because  $\forall \tilde{M}$  be IgPOS in  $D$ ,  $r^{-1}(\tilde{M})$  is IgPOS in  $E$ . But  $r$  is not Igsm, because  $r^{-1}(\{1, 3\}) = \{e, d\}$  is not IgOS in  $E$ .

**Example 3.5.** Let  $E = \{d, v, p, t\}$  with topology  $\mu = \{\dot{E}, \emptyset, \tilde{B}, \tilde{J}, \tilde{X}, \tilde{N}\}$ , where  $\tilde{B} = \langle e, \{d\}, \{v, p, t\} \rangle$ ,  $\tilde{J} = \langle e, \emptyset, \emptyset \rangle$ ,  $\tilde{X} = \langle e, \emptyset, \{v, p, t\} \rangle$ ,  $\tilde{N} = \langle e, \{d\}, \emptyset \rangle$ ,  $Y = \{1, 3, 5\}$  with topology  $\gamma = \{\dot{D}, \emptyset, \tilde{L}, \tilde{P}\}$ , where  $\tilde{L} = \langle d, \{1\}, \{3, 5\} \rangle$ ,  $P = \langle d, \{1\}, \emptyset \rangle$ . Let mapping  $r : (E, \mu) \rightarrow (D, \gamma)$  defined by  $r(\{d\}) = \{1\}$ ,  $r(\{v\}) = r(\{p\}) = \{3\}$ ,  $r(\{t\}) = \{5\}$ . Therefore  $r$  is IgSsm, because  $\forall \tilde{M}$  be IgSOS in  $D$ ,  $r^{-1}(\tilde{M})$  is IgSOS in  $E$ .

But  $r$  is not Ig $\alpha$ sm, because  $r^{-1}(\{3, 5\}) = \{p, v, t\}$  is not Ig $\alpha$ OS in  $E$ .

**Example 3.6.** Let  $E = \{i, j\}$  with topology  $\mu = \{\dot{X}, \emptyset, \tilde{M}, \tilde{N}\}$ , where  $\tilde{M} = \langle e, \{i\}, \{j\} \rangle$ ,  $\tilde{N} = \langle e, \{i\}, \emptyset \rangle$  and  $D = \{3, 4\}$  with topology  $\gamma = \{\dot{D}, \emptyset, \tilde{M}, \tilde{W}\}$ , where  $\tilde{M} = \langle x, \emptyset, \{3\} \rangle$ ,  $\tilde{W} = \langle x, \emptyset, \emptyset \rangle$ . Let a mapping  $r : (E, \mu) \rightarrow (D, \gamma)$  defined by as  $r(\{i\}) = \{3\}$ ,  $r(\{j\}) = \{4\}$ . So that  $r$  is Ig $\alpha$ sm, because  $\forall \tilde{M}$  be Ig $\alpha$ OS in  $D$ ,  $r^{-1}(\tilde{M})$  is

Ig $\alpha$ OS in  $E$ . But  $r$  is not Igsm, because  $r^{-1}(\{4\}) = \{j\}$  is not IgOS in  $E$ .

**Remark 3-7.** IgPsm and IgSsm is independent notions. The following two examples shows this two cases.

**Example 3.8.** Let  $E = \{w, r, i\}$  with topology  $\mu = \{\dot{E}, \emptyset, \tilde{K}, \tilde{S}, \tilde{G}, \tilde{Z}, \tilde{I}\}$ , where  $\tilde{K} = \langle e, \{w\}, \{r, i\} \rangle$ ,  $\tilde{S} = \langle e, \{w\}, \emptyset \rangle$ ,  $\tilde{G} = \langle e, \{w, r\}, \{i\} \rangle$ ,  $\tilde{Z} = \langle e, \{w\}, \{r\} \rangle$ ,  $\tilde{I} = \langle e, \{w, r\}, \emptyset \rangle$  and

$D = \{2, 4, 6\}$  with topology  $\gamma = \{\dot{D}, \emptyset, \tilde{T}, \tilde{H}, \tilde{C}, \tilde{F}, \tilde{R}\}$ , where  $\tilde{T} = \langle d, \{2\}, \{4, 6\} \rangle$ ,  $\tilde{H} = \langle d, \{2\}, \emptyset \rangle$ ,

$\tilde{C} = \langle d, \{2, 4\}, \{6\} \rangle$ ,  $\tilde{F} = \langle d, \{2\}, \{6\} \rangle$ ,  $\tilde{R} = \langle d, \{2, 4\}, \emptyset \rangle$ . Let a mapping  $r : (E, \mu) \rightarrow (D, \gamma)$  defined by  $r(\{w\}) = \{2\}$ ,  $r(\{r\}) = \{4\}$ ,  $r(\{i\}) = \{6\}$ . So  $r$  is IgPsm,

because  $\forall \tilde{M}$  be an IgPOS in  $D$ ,  $r^{-1}(\tilde{M})$  is IgPOS in  $E$ . But  $r$  is not IgSsm, because  $r^{-1}(\{4, 6\}) = \{w, i\}$  is not IgSOS in  $E$ .

**Example 3.9.** Let  $E = \{o, p, u\}$  with topology  $\mu = \{\dot{E}, \emptyset, \tilde{W}, \tilde{R}, \tilde{K}, \tilde{P}, \tilde{T}, \tilde{H}, \tilde{Z}\}$ , where

$\tilde{W} = \langle e, \{o\}, \{p, u\} \rangle$ ,  $\tilde{R} = \langle e, \{o\}, \{p\} \rangle$ ,  $\tilde{K} = \langle e, \{o\}, \emptyset \rangle$ ,  $\tilde{P} = \langle e, \{o, p\}, \emptyset \rangle$ ,  $\tilde{T} = \langle e, \emptyset, \emptyset \rangle$ ,

$\tilde{H} = \langle e, \emptyset, \{p, u\} \rangle$ ,  $\tilde{Z} = \langle e, \emptyset, \{p\} \rangle$  and  $D = \{1, 4, 7\}$  with topology  $\gamma = \{\dot{D}, \emptyset, \tilde{V}, \tilde{G}, \tilde{Q}\}$ , where  $\tilde{V} = \langle d, \{1\}, \{4, 7\} \rangle$ ,  $\tilde{G} = \langle d, \{1, 4\}, \emptyset \rangle$ ,  $\tilde{Q} = \langle d, \{1\}, \emptyset \rangle$ . Let a mapping  $r : (E, \mu) \rightarrow (D, \gamma)$

defined by  $r(\{o\}) = \{1\}$ ,  $r(\{p\}) = \{4\}$ ,  $r(\{u\}) = \{7\}$ . So  $r$  is IgSsm,

because for each  $\tilde{M}$  is IgSOS in  $D$ ,  $r^{-1}(\tilde{M})$  is IgSOS in  $E$ . But  $r$  is not IgPsm,

because  $r^{-1}(\{4, 7\}) = \{p, u\}$  is not IgPOS in  $E$ .

## References

[1] Zadeh, L. A., " Fuzzy sets", Information and control, 8, 338-353, 1965.  
[2] C.Chang, Fuzzy topological space, J.Math.Anal.Appl.24(1968),182-190  
[3] K. Atanassov, Intuitionistic fuzzy sets, VII ITKR's Session (Sofia, June 1983 Central Sci and Tech. Library) (V. Sgurev, ed.), Blug. Academy of Sciences, Sofia, 1984.  
[4] K. Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets and Systems, 20, (1986) , 87-96.

[5] D. Coker, " A note on intuitionistic sets and intuitionistic points" Turkish J. Math. 20, 1996 , 343-351.  
[6] D. Coker, An introduction to intuitionistic fuzzy topological space, Fuzzy Sets and Systems, 88 , 1997, 81-89.  
[7] D. Coker, D. An introduction to intuitionistic topological spaces, BUSEFAL 81, 51–56, 2000.  
[8] .Jeon, J. K., Jun, Y. B. and Park, J. H. (2005) " Intuitionistic fuzzy alpha-continuity and Intuitionistic fuzzy pre-continuity" I.J. of math., and Math. Sci. 19, pp. 3091-3101.

## حول تعميم بعض الاشكال الضعيفة للتطبيقات الفوقية في الفضاءات التبولوجية الحدسية

سامر رعد ياسين<sup>1</sup> ، هند فاضل عباس<sup>2</sup>

<sup>1</sup>قسم الرياضيات ، كلية علوم الحاسوب والرياضيات ، جامعة تكريت ، تكريت ، العراق

<sup>2</sup>متوسطة خالد ابن الوليد ، مديرية تربية صلاح الدين ، تكريت ، العراق

### الملخص

في هذا البحث قدمنا صفوف جديدة من التطبيقات اسميناها : intuitionistic generalized Pre supra mapping, intuitionistic generalized Semi supra mapping, intuitionistic generalized  $\alpha$ -supra mapping , intuitionistic generalized  $\beta$ -supra mapping ودرسنا بعض خواصها. واخيرا درسنا واستقصينا العلاقات بين هذه المفاهيم.