



GENERALIZED CONHARMONIC CURVATURE TENSOR OF NEARLY KAHLER MANIFOLD

Ali A. Shihab , Dhabia'a M. Ali

Department of Mathematics , College of Education for pure sciences, Tikrit University , Tikrit , Iraq

ARTICLE INFO.

Article history:

-Received: 12 / 9 / 2017

-Accepted: 30 / 10 / 2017

-Available online: / / 2018

Keywords: Generalized Conharmonic Curvature Tensor, Nearly Kahler Manifold.

Corresponding Author:

Name: Ali A. Shihab

E-mail: Ali.abd82@yahoo.com

Tel:

1- Introduction

Conformal transformations of Riemannian structures are the very significant object of study differential geometry, where this transformation which keeping the property of smooth harmonic function. It is known, that such transformations have tensor in variant so-called generalized conharmonic curvature tensor, The first practices were in 1960 by Koto [1] when he constructed a manifold that carried the properties of kahler manifold. In 1975, Cray [2] found an accurate method for forming new examples on practical kinds of manifold, which were based on the remarks of Koto.

In 1975 a great change was made on these studies by the Russian researcher Kirichenko, when he studied the almost Hermitian manifold by adjoined G-structure space in particular, he defined two tensors which were the structure and virtual tensors [3]. These tensors helped him to find the structure group of almost Hermitian manifold. Later on, the studies started to represent this method which allowed to study the geometrical structure characteristics for each class of almost Hermitian manifold. This study encouraged the researchers who were working with Kirichenko. In 1993, Banaru [4] succeeded in re-classifying the sixteen classes of almost Hermitian manifold by using the structure and Virtual tensors, which were named Kirichenko's tensors [5]. Among the sixteen classes of almost Hermitian manifold there are eight which are invariant under the conformal transformation metric.

Abstract

In this paper we study the relationship between tensor algebraic curvature tensor, and General conharmonic curvature tensor of Nearly Kahler manifold, i. e. it has a classical symmetry properties of the Riemann curvature tensor. Relapsing generalized Riemannian structure of certain classes of almost Hermitian manifold allows an additional symmetry properties of this tensor.

In this paper we investigated the generalized conharmonic curvature tensor of nearly kahler manifold.

2- Preliminaries

let M be a smooth manifold of dimension $2n$, $C^\infty(M)$ is algebra of smooth function on M ; $X(M)$ is the module of smooth vector fields on manifold of M ; $g = \langle \cdot, \cdot \rangle$ is Riemannian metrics, ∇ is Riemannian connection of the metrics g on M ; d is the operator of exterior differentiation. In the further all manifold, Tensor field, etc. objects are assumed smooth a class $C^\infty(M)$.

Definition 1.[6]: Almost Hermitian structure on a manifold (AH) M is the pair (J, g) , where J is an almost complex structure ($J^2 = -id$) on M , $g = \langle \cdot, \cdot \rangle$ is a (pseudo)Riemannian metric on M . In this case $\langle JX, JY \rangle = -\langle X, Y \rangle$, $\forall X, Y \in X(M)$.

Lemma 1.[7]: Every almost complex manifold there exist an almost Hermitian (AH)-structure. The endomorphism J is called a structural endomorphism, Manifold which is fixed on it an almost Hermitian structure is called an almost Hermitian manifold (AH) -manifold and denoted by $\{M, J, g = \langle \cdot, \cdot \rangle\}$ or simply M .

Definition2.[2]: Almost Hermitian structure (J, g) on manifold M is called;

1 -A Kahler (K) -structure, if $\nabla_X(J)Y = 0$; $X, Y \in X(M)$.

2 -A nearly Kahler (NK)-structure, if M satisfies $\nabla_X(J)Y + \nabla_Y(J)(X) = 0$; $X, Y \in X(M)$.

Lemma 2.[8]: An almost hermitian manifold successful :

- 1) $J^o \nabla_x(J)Y + \nabla_x(J)Y = 0$
- 2) $(\nabla_x(J)Y, Z) + (Y, \nabla_x(J)Z) = 0$

Theorem 3[8]: An Almost Hermitian structure (J, g) on a manifold M is nearly Kahler if and only if the following identities are fair:

- 1) $B(X, Y) = 0$
- 2) $C(X, Y) + C(Y, X) = 0$,
this structure is called Kahler iff $B=C=0$.

Definition 3.[8] : The set of tensors of type (r,1) are linear or antilinear in its argument and the sum components in the tensor T is called the spectrum of the tensor T, and the tensors themselves are called elements of the spectrum.

Definition: 4: Suppose (M, J, g) - Nk manifold. We retard alert, that curvature tensor general conharmonic was introduced by Ishii (1957) [9] as a tensor of type (4, 0) on n - dimensional Riemannian manifold, definite by the formulation:

$$K(HR)_{ijkl} = (HR)_{ijkl} - \frac{1}{2(n-1)} \{g_{ik} r(HR)_{jl} + g_{jl} r(HR)_{ik} - g_{il} r(HR)_{jk} - g_{jk} r(HR)_{il}\} \dots \dots (1)$$

Where (HR) is the general Riemann curvature tensor, r(HR) is general Ricci tensor. This tensor is invariant under general conharmonic transformation, i.e. with conformal transformations of space keeping a harmony of functions.

Consider properties tensor general conharmonic curvature $K(HR)$:

$$1- K(HR)(X, Y, Z, W) = (HR)(X, Y, Z, W) - 1/2(n-1) \{g(x,z) r(HR)(Y, W) + g(Y, W) r(HR)(X, Z) - g(X, W) r(HR)(Y, Z) - g(Y, Z) r(HR)(X, W)\} = -(HR)(Y, X, Z, W) + 1/2(n-1) \{g(X, Z) r(HR)(Y, W) + g(Y, W) r(HR)(X, Z) - g(X, W) r(HR)(Y, Z) - g(Y, Z) r(HR)(X, W)\} = -K(HR)(Y, X, Z, W)$$

The next ownerships are similarity proved

- 2- $K(HR)(X, Y, Z, W) = -K(HR)(X, Y, W, Z)$
- 3- $K(HR)(X, Y, Z, W) = K(HR)(Z, W, X, Y)$
- 4- $K(HR)(X, Y, Z, W) + K(HR)(X, Z, W, Y) + K(HR)(X, W, Y, Z) = 0$

So general conharmonic curvature tensor $K(HR)$ satisfies all the properties of algebraic curvature tensor :

- 1- $K(HR)(X, Y, Z, W) = -K(HR)(Y, X, Z, W)$
- 2- $K(HR)(X, Y, Z, W) = -K(HR)(X, Y, W, Z)$
- 3- $K(HR)(X, Y, Z, W) = K(HR)(Z, W, X, Y)$
- 4- $K(HR)(X, Y, Z, W) + K(HR)(X, Z, W, Y) + K(HR)(X, W, Y, Z) = 0, X, Y, Z, W \in X(M) \dots (2)$

Tensor generalized conharmonic curvature $K(HR)$ looks like :

$$K(HR)(X, Y)Z = (HR)(X, Y)Z - 1/2(n-1) \{r(HR)(Y, Z)X + r(HR)(X, Z)Y - g(Y, Z)QX - g(X, Z)QY\} \dots (3)$$

Where $g(QX, Y) = r(HR)(X, Y)$. By definition of aspectrum tensor, $K_3(HR)(X, Y)Z + K_4(HR)(X, Y)Z + K_5(HR)(X, Y)Z + K_6(HR)(X, Y)Z + K_7(HR)(X, Y)Z ; X, Y, Z \in X(M)$

It agrees Lemma 2: tensor $K_0(HR)(X, Y)Z$ as nonzero the component can have only components of the form

$\{K_0(HR)_{bcd}^a, K_0(HR)_{bcd}^{\hat{a}}\} = \{K(HR)_{bcd}^a, K(HR)_{bcd}^{\hat{a}}\}$ components of the from $\{K_1(HR)_{bcd}^a, K_1(HR)_{bcd}^{\hat{a}}\} = \{K(HR)_{bcd}^a, K(HR)_{bcd}^{\hat{a}}\}$; tensor $K_2(HR)(X, Y)Z$ - ingredients of the formulation $\{K_2(HR)_{bcd}^a, K_2(HR)_{bcd}^{\hat{a}}\} = \{K(HR)_{bcd}^a, K(HR)_{bcd}^{\hat{a}}\}$; tensor $K_3(HR)(X, Y)Z$ - ingredients of the form $\{K_3(HR)_{bcd}^a, K_3(HR)_{bcd}^{\hat{a}}\} = \{K(HR)_{bcd}^a, K(HR)_{bcd}^{\hat{a}}\}$; tensor $K_4(HR)(X, Y)Z$ - ingredients of the from $\{K_4(HR)_{bcd}^a, K_4(HR)_{bcd}^{\hat{a}}\} = \{K(HR)_{bcd}^a, K(HR)_{bcd}^{\hat{a}}\}$; tensor $K_5(HR)(X, Y)Z$ -ingredients of the form $\{K_5(HR)_{bcd}^a, K_5(HR)_{bcd}^{\hat{a}}\} = \{K(HR)_{bcd}^a, K(HR)_{bcd}^{\hat{a}}\}$; tensor $K_6(HR)(X, Y)Z$ - ingredients of the form $\{K_6(HR)_{bcd}^a, K_6(HR)_{bcd}^{\hat{a}}\} = \{K(HR)_{bcd}^a, K(HR)_{bcd}^{\hat{a}}\}$; tensor $K_7(HR)(X, Y)Z$ - ingredients of form $\{K_7(HR)_{bcd}^a, K_7(HR)_{bcd}^{\hat{a}}\} = \{K(HR)_{bcd}^a, K(HR)_{bcd}^{\hat{a}}\}$

Tensors $K_0(HR) = K_0(HR)(X, Y)Z, K_1(HR) = K_1(HR)(X, Y)Z, \dots, K_7(HR) = K_7(HR)(X, Y)Z$ we shall name the basic invariants conharmonic AH-manifold.

Now we introduces anew definition:-

Definition 5: AH-manifold for each $K_i(HR) = 0$, We shall name AH - manifold of class $K_i(HR)$, $i=0, 1, \dots, 7$.

Theorem2:

1) AH-manifold of class $K_0(HR)$ are characterized by identity

$$K(HR)(X, Y)Z - K(HR)(X, JY)JZ - K(HR)(JX, Y)JZ - K(HR)(JX, JY)Z - JK(HR)(X, Y)JZ - JK(HR)(JX, Y)Z - JK(HR)(JX, Y)Z + JK(HR)(JX, JY)JZ = 0; X, Y, Z \in X(M).$$

2) AH-manifold of class $K_1(HR)$ are characterized by identity

$$K(HR)(X, Y)Z + K(HR)(X, JY)JZ - K(HR)(JX, Y)JZ + K(HR)(JX, JY)Z + JK(HR)(X, Y)JZ - JK(HR)(X, JY)Z - JK(HR)(JX, Y)Z - JK(HR)(JX, JY)JZ ; X, Y, Z \in X(M).$$

3) AH-manifold of class $K_2(HR)$ are characterized by identity

$$K(HR)(X, Y)Z - K(HR)(X, JY)JZ + K(HR)(JX, Y)JZ + K(HR)(JX, JY)Z - JK(HR)(X, Y)JZ - JK(HR)(X, JY)Z + JK(HR)(JX, Y)Z - K(HR)(JX, JY)JZ = 0; X, Y, Z \in X(M).$$

4) AH-manifold of class $K_3(HR)$ are characterized by identity

$$K(HR)(X, Y)Z + K(HR)(X, JY)JZ + K(HR)(JX, Y)JZ - K(HR)(JX, JY)Z - JK(HR)(X, Y)JZ + JK(HR)(X, JY)Z + JK(HR)(JX, Y)Z + JK(HR)(JX, JY)JZ ; X, Y, Z \in X(M)$$

5) AH-manifold of class $K_4(HR)$ are characterized by identity

$$K(HR)(X, Y)Z + K(HR)(X, JY)JZ + K(HR)(JX, Y)JZ - K(HR)(JX, JY)Z + JK(HR)(X, Y)JZ - JK(HR)(X, JY)Z - JK(HR)(JX, Y)Z - JK(HR)(JX, JY)JZ = 0 ; X, Y, Z \in X(M).$$

6) AH-manifold of class $K_5(HR)$ are characterized by identity

$$K(HR)(X, Y)Z - K(HR)(X, JY)JZ + K(HR)(JX, Y)JZ + K(HR)(JX, JY)Z + JK(HR)(X, Y)JZ - JK(HR)(X, JY)Z - JK(HR)(JX, Y)Z - JK(HR)(JX, JY)JZ ; X, Y, Z \in X(M)$$

$K(HR)(X, Y)JZ + JK(HR)(X, JY)Z - JK(HR)(JX, Y)Z + JK(HR)(JX, JY)JZ = 0; X, Y, Z \in X(M)$.
7) AH-manifold of class $K_6(HR)$ are characterized by identity $K(HR)(X, Y)Z + K(HR)(X, JY)JZ - K(HR)(JX, Y)JZ + K(HR)(JZ, JY)Z + JK(HR)(X, Y)JZ - JK(HR)(X, JY)Z + JK(HR)(JX, Y)Z + JK(HR)(JX, JY)JZ = 0; X, Y, Z \in X(M)$.

8) AH-manifold of class $K_7(HR)$ are characterized by identity $K(HR)(X, Y)Z - K(HR)(X, JY)JZ - K(HR)(JX, Y)JZ - K(HR)(JX, JY)Z + JK(HR)(X, Y)JZ + JK(HR)(X, JY)Z + JK(HR)(JX, Y)Z - JK(HR)(JX, JY)JZ = 0; X, Y, Z \in X(M)$.

Proof:

The manifold of class $K_0(HR)$ is characterized by a condition $K_0(HR)^a_{bcd} = 0$, or $K^a_{bcd} = 0$, i.e. $[k(HR)(\varepsilon_c, \varepsilon_d)\varepsilon_b]^a \varepsilon_a = 0$. As σ – a projector on D_j^{-1} , that $\sigma \circ \{K(HR)(\sigma X, \sigma Y)\sigma Z\} = 0$, i.e. $(id - \sqrt{-1}j)\{K(HR)(X - \sqrt{-1}jX, Y - \sqrt{-1}jY)(Z - \sqrt{-1}jZ)\} = 0$. Removing the brackets we shall receive:

$$K(HR)(X, Y)Z - K(HR)(X, JY)JZ - K(HR)(JX, Y)JZ - K(HR)(JX, JY)Z - JK(HR)(X, Y)JZ - JK(HR)(X, JY)Z - JK(HR)(JX, Y)Z + JK(HR)(JX, JY)JZ - \sqrt{-1}\{K(HR)(X, Y)JZ + K(HR)(X, JY)Z + K(HR)(JX, Y)Z - K(HR)(JX, JY)JZ\} - \{JK(HR)(X, Y)Z - JK(HR)(X, JY)JZ - JK(HR)(JX, Y)Z - JK(HR)(JX, JY)JZ\} = 0,$$

i.e.

$$1) K(HR)(X, Y)Z - K(HR)(X, JY)JZ - K(HR)(JX, Y)JZ - K(HR)(JX, JY)Z - JK(HR)(X, Y)JZ - JK(HR)(X, JY)Z - JK(HR)(JX, Y)Z + JK(HR)(JX, JY)JZ = 0$$

Thus AH-manifold of class $K_0(HR)$ are characterized by identity

$$2) K(HR)(X, Y)JZ + K(HR)(X, JY)Z + K(HR)(JX, Y)Z - K(HR)(JX, JY)JZ + JK(HR)(X, Y)Z - JK(X, JY)JZ - JK(HR)(JX, Y)Z - JK(HR)(JX, JY)Z = 0, X, Y, Z \in X(M)$$

These equality are equivalent, the second equality turns out from the first replacement Z on JZ .

$$K(HR)(X, Y)Z - K(HR)(X, JY)JZ - K(HR)(JX, Y)JZ - K(HR)(JX, JY)Z - JK(HR)(X, Y)JZ - JK(HR)(X, JY)Z - JK(HR)(JX, Y)Z + JK(HR)(JX, JY)JZ = 0, X, Y, Z \in X(M)$$

Similarly, considering AH-manifold of class $K_1(HR) - K_7(HR)$, we shall receive the following theorem.

Theorem 3.[3]: Components of Riemannian curvature tensor of NK-manifold in the adjoint G-structurespace are award as follows :

$$R^a_{bcd} = R^a_{b\hat{c}\hat{d}} = R^{\hat{a}}_{bcd} = R^{\hat{a}}_{b\hat{c}\hat{d}} = R^{\hat{a}}_{bcd} = R^{\hat{a}}_{bc\hat{d}} = R^{\hat{a}}_{b\hat{c}\hat{d}} = R^{\hat{a}}_{b\hat{c}\hat{d}} = R^{\hat{a}}_{b\hat{c}\hat{d}} = 0$$

- 1) $R^a_{bcd} = B^{adh}B_{hbc} + A_{bc}^a$
- 2) $R^a_{b\hat{c}\hat{d}} = B^{abc}B_{bdh} - A_{bd}^a$
- 3) $R^{\hat{a}}_{bcd} = B^{bdh}B_{ahc} - A_{ac}^{\hat{a}}$
- 4) $R^{\hat{a}}_{b\hat{c}\hat{d}} = B^{bch}B_{adh} + A_{ad}^{\hat{a}}$
- 5) $R^{\hat{a}}_{b\hat{c}\hat{d}} = 2B^{dch}B_{abh}$
- 6) $R^{\hat{a}}_{b\hat{c}\hat{d}} = 2B^{abh}B_{dch}$

Definition 6.[10]: Ageneralized Riemannian curvature tensor on AH-manifold M is a tensor of type (4,0) which is defined as following formulation:

$$(HR)(X, Y, Z, W) = \frac{1}{16} \{3[R(X, Y, Z, W) + R(JX, JY, Z, W) + R(X, Y, JZ, JW) + R(JX, JY, JZ, JW] - R(X, Z, JW, JY) - R(JX, JZ, W, Y) - R(X, W, JY, JZ) - R(JX, JW, Y, Z) + R(JX, Z, JW, Y) + R(X, JZ, W, JY) + R(JX, W, Y, JZ) + R(X, JW, JY, Z)\}$$

Where $R(X, Y, Z, W)$ is Riemannian curvature tensor, $X, Y, Z, W \in T_p(M)$ and satisfies the following properties:

- 1) $(HR)(X, Y, Z, W) = -(HR)(Y, X, Z, W) = -(HR)(X, Y, W, Z)$;
- 2) $(HR)(X, Y, Z, W) = (HR)(Z, W, X, Y)$;
- 3) $(HR)(X, Y, Z, W) + (HR)(X, Z, W, Y) + (HR)(X, W, Y, Z) = 0$;
- 4) $(HR)(X, JX, JX, X) = R(X, JX, JX, X)$

The components of the generalized Riemannian curvature tensor (HR) of NK-manifold are given by the next theorem.

Theorem 4.[6]: The components of the generalized Riemannian curvature by the next formulation :

- 1) $(HR)_{abcd} = -A_{bd}^{ac}$
- 2) $(HR)_{\hat{a}bc\hat{d}} = A_{bc}^{\hat{a}d}$

And the others are conjugate to the above components or equal to zero.

Definition 7.[11]: A tensor of type (2,0) which is defined as $r(HR)_{ij} = (HR)_{ijk}$ is called a generalized Ricci tensor.

Theorem 5.[12]: The components of generalized Ricci tensor of NK-manifold in the adjoint G-structure space are given as the following form:

$$R(HR)_{ab} = -A_{bc}^{ac}$$

And the others are conjugate to the above component or equal to zero.

Definition 8.[12]: A generalized scalar curvature tensor is denoted by $K(HR)$ and defined as :

$$K(HR) = g_{ij} r(HR)_{ij}$$

Theorem 6.[12]: The component of generalized scalar curvature tensor of NK-manifold in the adjoint G-structure space is given as the following form :

$$K(HR) = -2A_{ac}^{ac}$$

Theorem 7: The components of the generalized conharmonic curvature tensor of NK-manifolds in the adjointG-structure are given as the following form:

$$K(HR)_{ijkl} = R(HR)_{ijkl} - \frac{1}{2(n-1)} \{g_{ik} r(HR)_{jl} + g_{jl} r(HR)_{ik} - g_{il} r(HR)_{jk} - g_{jk} r(HR)_{il}\}$$

$$1- K(HR)_{\hat{a}b\hat{c}\hat{d}} = -A_{bd}^{ac} - 1/2(n-1) \{ \delta_a^{\hat{a}} r(HR)_{\hat{c}\hat{b}} + \delta_b^{\hat{a}} r(HR)_{\hat{a}\hat{d}} \}$$

$$2- K(HR)_{\hat{a}bc\hat{d}} = A_{bc}^{\hat{a}d} + 1/2(n-1) \{ \delta_c^{\hat{a}} r(HR)_{b\hat{a}} + \delta_b^{\hat{a}} r(HR)_{\hat{a}\hat{c}} \}$$

$$3- K(HR)_{\hat{a}\hat{b}cd} = 1/2(n-1) \{ \delta_c^{\hat{a}} r(HR)_{\hat{b}c} + \delta_d^{\hat{a}} r(HR)_{\hat{a}\hat{c}} - \delta_a^{\hat{a}} r(HR)_{\hat{b}c} - \delta_c^{\hat{b}} r(HR)_{\hat{a}\hat{d}} \}$$

And other components Generalized conharmonic curvature tensor for NK are equal to zero.

Proof:

1- Let $i=a, j=b, k=c$ and $l=d$

$$K(HR)_{abcd} = (HR)_{abcd} - \frac{1}{2(n-1)} \{ g_{ac} r(HR)_{bd} + g_{bd} r(HR)_{ac} - g_{ad} r(HR)_{bc} - g_{bc} r(HR)_{ad} \}.$$

$$k_{abcd} = 0 - \frac{1}{2(n-1)} \{ (0)(0) + (0)(0) - (0)(0) - (0)(0) \} = 0$$

$$k(HR)_{abcd} = 0$$

2-Let $i=\hat{a}$, $j= b$, $k =c$ and $l=d$

$$K(HR)_{\hat{a}bcd} = (HR)_{\hat{a}bcd} - \frac{1}{2(n-1)} \{ g_{\hat{a}c} r(HR)_{bd} + g_{bd} r(HR)_{\hat{a}c} - g_{\hat{a}d} r(HR)_{bc} - g_{bc} r(HR)_{\hat{a}d} \}$$

$$k(HR)_{\hat{a}bcd} = 0 - \frac{1}{2(n-1)} \{ \delta_c^a(0) + (0)(-A_{ck}^{ak}) - \delta_d^a(0) - (0)A_{dk}^{ak} - \frac{1}{2(n-1)} \{0\}$$

$$k(HR)_{\hat{a}bcd} = 0$$

3-Let $i=a$, $j=\hat{b}$, $k =c$ and $l=d$

$$K(HR)_{a\hat{b}cd} = (HR)_{a\hat{b}cd} - \frac{1}{2(n-1)} \{ g_{\hat{a}c} r(HR)_{\hat{b}d} + g_{\hat{b}d} r(HR)_{\hat{a}c} - g_{\hat{a}d} r(HR)_{\hat{b}c} - g_{\hat{b}c} r(HR)_{\hat{a}d} \}$$

$$k(HR)_{a\hat{b}cd} = 0 - \frac{1}{2(n-1)} \{ \delta_c^a(0) + (0)(-A_{ck}^{ak}) - \delta_d^a(0) - (0)(-A_{dk}^{ak}) \}$$

$$= -\frac{1}{2(n-1)} \{0\}$$

$$K(HR)_{a\hat{b}cd} = 0$$

4-Let $i=a$, $j= b$, $k=\hat{c}$ and $l=d$

$$K(HR)_{ab\hat{c}d} = (HR)_{ab\hat{c}d} - \frac{1}{2(n-1)} \{ g_{\hat{a}c} r(HR)_{bd} + g_{bd} r(HR)_{\hat{a}c} - g_{\hat{a}d} r(HR)_{bc} - g_{bc} r(HR)_{\hat{a}d} \}$$

$$k(HR)_{ab\hat{c}d} = 0 - \frac{1}{2(n-1)} \{ \delta_c^a(0) + (0)(-A_{ck}^{ak}) - (0)(-A_{bk}^{ck}) - \delta_b^c(0) \}$$

$$= -\frac{1}{2(n-1)} \{0\}$$

$$K(HR)_{ab\hat{c}d} = 0$$

5-Let $i=a$, $j= b$, $k=c$ and $l=\hat{d}$

$$K(HR)_{abc\hat{d}} = (HR)_{abc\hat{d}} - \frac{1}{2(n-1)} \{ g_{ac} r(HR)_{b\hat{d}} + g_{b\hat{d}} r(HR)_{ac} - g_{a\hat{d}} r(HR)_{bc} - g_{bc} r(HR)_{a\hat{d}} \}$$

References

[1] Koto, S. (1960). Some Theorems on Almost Kahlerian Spaces. Journal Math Soc. Japan, **12**:422-433.
 [2] Gray, A. (1970). Nearly Kahler Manifold. J. Diff. Geom. **4(3)**:283-309.
 [3] Kirichenko, V. F. (1975). New Results of K-space Theory. ph. D. thesis, Moscow state University of Moscow.
 [4] Banaru, M. (1993). Hermitian geometry of six-dimensional submanifold of cayley's algebra. ph. D. thesis, Moscow state pedagogical university, Moscow.
 [5] Banaru, M. (2001). A new characterization of the Gray-Hervalla classes of almost Hermitian manifold. 8th International conference on differential geometry and its applications, 27-31 Augus 2001, Opava-Czech Republic.
 [6] Shihab, A.A. (2011). On the geometry of conharmonic curvature tensor of nearly kahler manifold. Jounal of Basrah Researches, **37(4)**, 39-48.

$$k(HR)_{abc\hat{d}} = 0 - \frac{1}{2(n-1)} \{ (0)(-A_{bk}^{dk}) + \delta_b^d(0) - \delta_a^d(0) - (0)(-A_a^d) \}$$

$$= -\frac{1}{2(n-1)} \{0\}$$

$$K(HR)_{abc\hat{d}} = 0$$

6-Let $i=\hat{a}$, $j=\hat{b}$, $k=c$ and $l=d$

$$K(HR)_{\hat{a}\hat{b}cd} = (HR)_{\hat{a}\hat{b}cd} - \frac{1}{2(n-1)} \{ (HR)_{\hat{a}c} r(HR)_{\hat{b}d} + g_{\hat{b}c} r(HR)_{\hat{a}c} - g_{\hat{a}d} r(HR)_{\hat{b}c} - g_{bc} r(HR)_{\hat{a}d} \} = 0 - \frac{1}{2(n-1)} \{ \delta_c^a(-A_{dk}^{bk}) + \delta_d^b(-A_{ck}^{ak}) - \delta_d^a(-A_{ck}^{bk}) - \delta_c^b(-A_{dk}^{ak}) \}$$

$$K(HR)_{\hat{a}\hat{b}cd} = \frac{1}{2(n-1)} \{ \delta_c^a A_{dk}^{bk} + \delta_d^b A_{ck}^{ak} - \delta_d^a A_{ck}^{bk} - \delta_c^b A_{dk}^{ak} \}$$

7-Let $i=\hat{a}$, $j=b$, $k=\hat{c}$ and $l=d$

$$K(HR)_{\hat{a}b\hat{c}d} = (HR)_{\hat{a}b\hat{c}d} - \frac{1}{2(n-1)} \{ g_{\hat{a}c} r(HR)_{bd} + g_{bd} r(HR)_{\hat{a}c} - g_{\hat{a}d} r(HR)_{b\hat{c}} \} =$$

$$-A_{bd}^{ac} - \frac{1}{2(n-1)} \{0 + 0 - \delta_d^a(-A_{bk}^{ck}) - \delta_b^c(-A_{dk}^{ak}) \}$$

$$K(HR)_{\hat{a}b\hat{c}d} = -A_{bd}^{ac} - \frac{1}{2(n-1)} \{ \delta_d^a A_{bk}^{ck} + \delta_b^c A_{dk}^{ak} \}$$

8-Let $i=\hat{a}$, $j=b$, $k=c$ and $l=\hat{d}$

$$K(HR)_{\hat{a}bc\hat{d}} = (HR)_{\hat{a}bc\hat{d}} - \frac{1}{2(n-1)} \{ g_{\hat{a}c} r(HR)_{b\hat{d}} + g_{\hat{a}d} r(HR)_{\hat{a}c} - g_{\hat{a}d} r(HR)_{bc} - g_{bc} r(HR)_{\hat{a}d} \} = A_{bc}^{ad} - \frac{1}{2(n-1)} \{ \delta_c^a(-A_{bk}^{dk}) + \delta_b^d(-A_{ck}^{ak}) - 0 - 0 \}$$

$$K(HR)_{\hat{a}bc\hat{d}} = A_{bc}^{ad} + \frac{1}{2(n-1)} \{ \delta_c^a A_{bk}^{dk} + \delta_b^d A_{ck}^{ak} \} \blacksquare$$

[7] Kirichenko, F.V.; Rustanov A.R. and Shihab A. (2011). Geometry curvature tensor conharmonic almost Hermitian manifold. // Mat. Notes, t.90,No-1.
 [8] Kirichenko V.F. (2003). Differential-Geometrical Structure on Manifold. -M., MSPU, 495.
 [9] Ishii, Y. (1957). On Conharmonic Transformations Tensor. **7(2)**, 73-80.
 [10] Gan, G. (1978). Characteristic of some classes of almost Harmitin manifolds . Serdica, **4**, 19-23.
 [11] Mileva, p. (2003). Locally Conformally Kahler Manifolds of Constant tupe and J-invariant Corvature Tensor. Facta universitatis, Series: Mechanacs, Automatic control and Robotics , **3(14)**, 791.804.
 [12] Saadan, M.J. (2011). Complex Concircular Curvature Tensor of Certain Classes of Almost Hermitian Manifold. (Master thesis). University of Basrah.

تنزr الانحناء الكونهورمني المعمم لمنطوي كوهلر التقريبي

علي عبد المجيد شهاب ، ضبياء محمود علي حسين

قسم الرياضيات ، كلية التربية للعلوم الصرفة ، جامعة تكريت ، تكريت ، العراق

الملخص

في هذه البحث تم دراسة العلاقة بين تنزr الانحناء الجبري، وتنزr الانحناء الكونهورمني المعمم لمنطوي كوهلر التقريبي، حيث ان هذا التنزr يمتلك خصائص التناظر الكلاسيكي لتنزr الانحناء الريماني، مع احتساب مركبات تنزr الانحناء الكونهورمني المعمم في بعض أصناف المنطوي الهرميتي التقريبي إضافة الى دراسة خصائص التناظر لهذا التنزr.