



## Prevalence of *Toxoplasma gondii*, *Cytomegalovirus* and *Rubella virus* among aborted women in Samarra city

Sinai Najy Muhsin AL-doury

Collage of Dentistry, University of Tikrit , Tikrit , Iraq

### ARTICLE INFO.

#### Article history:

-Received: 18 / 5 / 2017

-Accepted: 28 / 1 / 2018

-Available online: / / 2018

**Keywords:** *Toxoplasma gondii*, *Cytomegalovirus*, *Rubella*, aborted, women, Samarra.

#### Corresponding Author:

Name: Sinai najy Muhsin

E-mail: [suinaldoury@gmail.com](mailto:suinaldoury@gmail.com)

Tel:

### Abstract

The aim of this study is to determine seroprevalence of *Toxoplasma gondii*, CMV and *Rubella virus* for 158 aborted women in Samarra General Hospital through period from January to December 2016. The prevalence of *T. gondii* was 14.6%, CMV 10.8%, *Rubella virus* 8.2% and common Toxo. & CMV was 9.5%. Also, the present study showed that infection with each of *T. gondii*, CMV and *Rubella* were highest (24.2%, 15.2%, 15.2% respectively) in age group 21-30 years old, while common infection with Toxo. & CMV was highest 11.8% in age group 14-20 years old. The seroprevalence results were for anti-*Toxo*. IgM was 7%, IgG was 3.2%, and IgM & IgG was 4.4%, anti-CMV IgM was 5.7%, IgG was 1.9%, and IgG & IgM was 3.2% and anti-*Rubella* IgM 3.8%, IgG 1.9%, and IgG & IgM was 2.5%. While the seroprevalence of both *Toxo*. & CMV was anti-*Toxo*. IgG & anti-CMV IgM was 0.63%, anti-*Toxo*. IgM & anti-CMV IgM was 2.5%, anti-*Toxo*. IgG & anti-CMV IgG & IgM was 1.9%, anti-*Toxo*. IgM & anti-CMV IgG & IgM was 3.2% and anti - *Toxo*. IgM & anti-CMV IgG was 1.3%.

### Introduction

Abortion is an elimination of fetus or embryo out women uterus. An abortion includes purposely (induced abortion) and may be caused spontaneously (miscarriage)[1] which causes emotional distress for parents[2].

many clinical types of abortion includes Complete, incomplete, threatened, missed, septic and non septic abortion. As well as many factors has been caused spontaneously. Abortion included uterine and genetic abnormalities, pollutants of environment, endometriosis, psycho-genetic factors[3]. For infectious agents such as *cytomegalovirus* and *Toxoplasma*, it is believed that 10-15% of miscarriage caused by those agents[4].

*Toxoplasma gondii* is one of intracellular widespread parasite[5] which infects human accidentally by ingestion of food water or soil contained oocysts of the parasite or by eating raw \undercooked meat containing cysts of *T. gondii*[6]. Human *Cytomegalovirus* is one of congenital malformation causes in both progressing and non progressing countries which responsible for asymptomatic infection to severe fetal damage[7].

Generally, infection of *Rubella* is an asymptotically childhood disease, but through the

pregnancy first trimester it responsible for congenital defects or fetal death congenital rubella syndrome (CRS)[8].

The chief aim of the current study is detection of antibodies to *Toxoplasma gondii*, *Cytomegalovirus*, *Rubella virus* and association infection of both *Toxo*. & CMV. Also, determine the potential role of those infectious agents in the incidence of abortion in aborted women attending General hospital in Samarra city in Salah al-Din province.

### Materials and methods

The study includes 158 aborted women attending Samarra General Hospital through the period from January to December 2016 to determine prevalence of antibody to *Toxoplasma gondii*, *Cytomegalovirus* and *Rubella virus*. Blood samples were examined for all aborted women to detection the presence of IgM & IgG to infectious agents. The ages of study groups ranged from 14-46 years old, Enzyme Linked Immunosorbent Assay (ELISA) were used to detect specific antibodies to *Toxo*., CMV and *Rubella* (from Biocheck, CA, USA), for association infection of *Toxo*. & CMV it was used (ELISA) Kit (Acon for sight(Germany)).

### Results

A total 158 aborted women were screened in Samarra General Hospital and the results were as follows:- The percentage of *Toxo. gondii* was 14.6%, CMV was 10.8%, *Rubella virus* was 8.2% and common *Toxo. & CMV* were 9.5%. Table (1).

**Table (1):- The prevalence of infectious agents among 158 aborted women**

infectious agent	No.	%
<i>Toxoplasma gondii</i>	23	14.6
Cytomegalovirus	17	10.8
<i>Rubella virus</i>	13	8.2
<i>Toxoplasma gondii</i> & Cytomegalovirus	15	9.5
Total	68	43.03

**Table (2):- The prevalence of infection according to age groups.**

Ages (years)	No. of tested women	infected women with <i>T. gondii</i>		infected women with CMV		infected women with <i>Rubella virus</i>		Infected women with <i>Toxo. &amp; CMV</i>	
		No.	%	No.	%	No.	%	No.	%
14-20	34	2	5.9	3	8.8	1	2.9	4	11.8
21-30	66	16	24.2	10	15.2	10	15.2	7	10.6
31-46	58	5	8.6	4	6.9	2	3.4	4	6.9
Total	158	23	14.6	17	10.8	13	8.2	15	9.5

The seroprevalence of anti-infectious agents immunoglobulins were:- for anti-*Toxo. IgM* was 7%, anti-*Toxo. IgG* was 3.2% and anti-*Toxo. IgM & IgG* were 4.4%. For anti-CMV IgM was 5.7%, anti-CMV IgG was 1.9% and anti-CMV IgM & IgG were 3.2%. For anti-*Rubella IgM* was 3.8%, anti-*Rubella IgG* was 1.9% and anti-*Rubella IgM & IgG* were 2.5%. Table (3).

**Table (3):- The seroprevalence of anti-Infectious agents immunoglobulins among 158 aborted women.**

Infectious agents	IgM		IgG		IgM & IgG	
	No.	%	No.	%	No.	%
<i>Toxoplasma gondii</i>	11	7	5	3.2	7	4.4
Cytomegalovirus	9	5.7	3	1.9	5	3.2
<i>Rubella virus</i>	6	3.8	3	1.9	4	2.5

The seroprevalences of both anti-*Toxo. & anti-CMV* were anti-*Toxo. IgG & anti-CMV IgM* were 0.63%, anti-*Toxo. IgM & anti-CMV IgM* were 2.5%, anti-*Toxo. IgG & anti-CMV IgG & IgM* were 1.9%, anti-*Toxo. IgM & anti-CMV IgG & IgM* were 3.2% and anti-*Toxo. IgM & anti-CMV IgG* were 1.3%. Table (4).

**Table (4):- The seroprevalence of association infection of *Toxo. & CMV*.**

Immunoglobulins	<i>Toxoplasma gondii</i> & Cytomegalovirus	
	No.	%
<i>Toxo IgG &amp; CMV IgM</i>	1	0.63
<i>Toxo. IgM &amp; CMV IgM</i>	4	2.5
<i>Toxo. IgG &amp; CMV (IgG &amp; IgM)</i>	3	1.9
<i>Toxo. IgM &amp; CMV (IgG &amp; IgM)</i>	5	3.2
<i>Toxo. IgM &amp; CMV IgG</i>	2	1.3

## Discussion

*Toxoplasma*, CMV and *Rubella* are important infectious agents causing infections which be mild or asymptomatic in the mother. previous agents may result in dangerous congenital abnormalities such as

The prevalence according to age groups in this study was:- for *Toxo.* was 5.9% in age group 14-20 years old, 24.2% in age group 21-30 years old and 8.6% in age group 31-46 years old.

For CMV was 8.8% in age group 14-20 years old, 15.2% in age group 21-30 years old and 6.9% in age group 31-46 years old. For *Rubella virus* was 2.9% in age group 14-20 years old, 15.2% in age group 21-30 years old and 3.4% in age group 31-46 years old. While the prevalence of association infection of *Toxo. & CMV* was 11.8% in age group 14-20 years old, 10.6% in age group 21-30 years old and 6.9% in age group 31-46 years old. Table (2).

intra-uterine growth, retardation, and even fetus death[1].

This study found that infection of *T. gondii* was 14.6%, it is differ from a study conducted in Al-Baiji[9], Najaf [10] and in Waste [11]. While the infection percentage of CMV was 10.8%, *Rubella virus* was 8.2% and common *Toxo. & CMV* were 9.5% respectively, it is differ from those reported in Waste[11], and Baghdad[12].

Add to the present study revealed that infection with *T. gondii*, CMV and Rubella were highest 24.2%, 15.2%, 15.2% respectively in age group 21-30 years old, while common infection with *Toxo. & CMV* was highest 11.8% in age group 14-20 years old.

Prevalence rate of infectious agents in this study probably due to the lack of presence of vaccination for *T. gondii* and CMV[9]. lack of drinking water chlorination in area of our study, increased number of stray animals especially in resent years may also contribute to the high rates and may be poverty about simple protective measures such as flushing all vegetables and fruits , hand washing , In addition to that, most of the families in the study area still raise cattle.

If a pregnant woman receives her exposure to *T. gondii* while the first three months of her pregnancy, the fetus will be at risk of abortion or congenital problems such as hydrocephalus, microcephalus and mental retardation. Being the parasite in uterus leads either chronic endometritis which interfere with fertilized egg or transferring through placenta to the fetus causing death[13].

CMV infections are mightly associated with popular life style, bad hygienic conditions[14]. However, we didn't collect data on *Rubella virus* vaccination from the women, so probably those aborted woman unvaccinated to prevent congenital rubella syndrome. women who contact with these viruses during the first three months of pregnancy will be affected. The

viruses invade the placenta and fetus during pregnancy causing birth to child with serious congenital problems such as blindness and deafness [15].

The seroprevalence of anti-infectious agents immunoglobulins in this study were for anti-*Toxo.* IgM was 7%, anti-*Toxo.* IgG was 3.2% and anti-*Toxo.* IgM & IgG were 4.4%. For anti-CMV IgM was 5.7%, anti-CMV IgG was 1.9% and anti-CMV IgM & IgG were 3.2%. For anti- *Rupella* IgM was 3.8%, anti- *Rupella* IgG was 1.9% and anti- *Rupella* IgM & IgG were 2.5%. Our previous results differ from that study conducted by Sadik *et. al.*[16] and those reported in each of Thiqr[17], Baghdad[12] and Kirkuk[18].

Also the seroprevalences of both anti-*Toxo.* & anti-CMV were for anti- *Toxo.* IgG & anti-CMV IgM were 0.63, anti-*Toxo.* IgM & anti-CMV IgM were 2.5%, anti-*Toxo.* IgG & anti-CMV IgG & IgM were 1.9%, anti-*Toxo.* IgM & anti-CMV IgG & IgM were 3.2% and anti-*Toxo.* IgM & anti-CMV IgG were 1.3%. those previous results differ from those reported in AL-Diwaniyah[19] and Thiqr[17]. The differences between our study results and these of previous studies may be due to the sensitivity differences between the serological tests, different numbers of tested women, the variations regions even within the same country.

*T. gondii* is the causative of toxoplasmosis, which its prevalence rates in the world may be changeable according to many factors, such nutritional habits (unwashing hands after contact with raw meat or contact with cats pets and consuming raw vegetables), lifestyle specially most of the families in the study area still raise cattle and Lack of knowledge of the parasite and its prevalence methods and the risk of the pregnant woman infection by awareness

## References

- 1- Stuart G and Grimes DA. (2010) Terminology of abortion. *J. Contra.*, **81** (2) pp.6- 93.
- 2- Khalida AI, Maysara SK and Dhamra WA. (2012) the seroprevalence study of cytomegalovirus between infected aborted women. *J. The Iraqi postgradu. Med.*, **11**(1) pp. 123.
- 3- Zuhara KF, Sekaran K and Sebastian D. (2008) effect of TORCH in first trimester miscarriage in the Malabar \ Kerala. *African J. Microbio. Res.*, **2** pp. 56-59.
- 4- Aggarwal A, Chopra S and Arora U. (2004) Prevalence of IgM to Toxoplasma, Rubella and Cytomegalovirus Infections. *J. K. scien.*, **6** (4) pp.190-192.
- 5- Koliou M, Detsika M, Sifakis S, Antoniou M and Messaritakis I. (2008) study of Geno-types of Toxoplasmosis in Pregnant Women & Patients from Crete and Cyprus. *Americ. Socie. Of Tropic. Medic. And Hyg.*, **79**(2) pp. 205-209.
- 6- Les TJ, Rosso F, Villalobos C, Chaves AJ, Agudelo A, Messa A, Remington SJ, Tunubala AG and Montoya GJ. (2008) Prevalence of

programs and unavailability of vaccine against the parasite[20].

The prevalence of CMV infections are associated with poor hygienic, low socio-economic level and crowded living conditions[21] specially The lifestyle of families living in large numbers is common in the society in which our study was conducted. embryonic CMV infection may happen following both primary and recurrent infection and reinfection with a new strain of the virus or latent infection reactivation can cause infections even in the being of IgG levels[20]. Risk for congenital Rubella should be done meticulously, even the rate of positive anti-Rubella IgM antibodies to be low[22]. Although successful rubella vaccine have been implemented in Iraq, the World Health Organization showed that Rubella is present in our country[23] and Majeed *et. al.*,[24] showed that there is an reduce of rubella immune which may be due to disturbance of the vaccination program in the time from 1992 to 2016, in addition to waning of vaccine stimulated immunity among women whom had received only one dose of Rubella vaccine.

A positive result of IgM for each of *T. gondii*, *Rubella*, and CMV may not always indicate a new primary infection. False-positive IgM results may occur in patients with autoimmune diseases due to the presence of rheumatoid factor and antinuclear antibodies or in patients recently infected with other viral pathogens due to heterotypic IgM antibody reactivity[25] specially we do not provide data about whether or not these women infected with another disease or how many times they have aborted.

In conclusion, more studies about common infection of both *T. gondii* and CMV should be conducted and their relationship to abortion.

Toxoplasmosis between Pregnant Women in Cali, Colombia/South America. *Americ. Soci. Of Trop. Medi. And Hyg.*, **78**(3) pp. 504-508.

7- Demmler GJ. (1991) Infectious diseases society of America and centers for disease control: Summary of a workshop on surveillance for congenital cytomegalovirus disease. *Rev. Infect. Dis.*, **13**(3) pp.15-29.

8- Abdul-Muhymen N, Al-Saadie M and Abdul-Karim ET. (2009) Chlamydia & rubella antibodies in women with full-term deliveries and women with abortion in Baghdad. *East. Medi. Heal. J.*, **15** (6) pp. 1407-1411.

9- Al-Jebouri M, Al-Janabi M and Ismail H. (2013) The prevalence of toxoplasmosis among female patients in Al-Hawija and Al-Baiji Districts in Iraq. *Open J. of Epidemio.*, **3** pp. 85-88.

10-AL- kalaby RF, AL-Fatlawi SN and Sultan BA. (2016) Relationship between abortion & Toxo. gondii in aborted women in Najaf province. *Scien. J. of Kerbala Univ.*, **14** (1) pp. 177-185.

- 11- Jasim M, Ali AI and Majeed HA. (2011) serological diagnosis of TORCH in both aborted and non aborted women in waste. *Tik. Med. J.*, **17**(2) pp. 141-147.
- 12- AL-Hindawi N Gh and AL-Shanawi FA. (2015) roprevalence of CMV, Toxo. gondii in aborted women in Baghdad\Iraq. *Iraqi J. of sciences.*, **56**(1) pp.694-655.
- 13- Meral T, Aysun K, Yusuf P, and Yasemin IB (2011) Evaluation of, *T. gondii* , cytomegalovirus & rubella seroprevalences between pregnant women in Denizli. *Turk. J. Med. Sci.*, **41** (1) pp. 159-164.
- 14- Vauloup-Fellous C, Picone O, Parent CI, Senat MV, Frydman R and Cordier AG.(2009) A two-years study on CMV infection during pregnancy in a French hospital. *BJOG.*, **116** PP. 818-823.
- 15- Robson J, Coulter C and Wood R. (1999) Rubella in pregnancy. *Commu. Disea. Intell. J.*, **23** pp. 93-96.
- 16- Sadik MS, Jamil K, Patil C and Fatima H. (2012) TORCH study in patients with bad obstetric history. *Bio. & Med.J.*, **4** (2) pp. 95-101.
- 17- Hadi NJ.(2011) Seroprevalence of CMV, Toxo. gondii & Rubella virus between aborted women in Thiqr. *J. of college of education for pure science.*, **5**(1) pp.3-9.
- 18- Hassan HMM, AL-Samarai AM, AL-Jumaili ZKM and AL-Obaidi AH. (2014) Association between CMV & bad obstetric outcomes in women at Kirkuk. *Internation. J. of Publ. Heal. Science.*, **3**(1) pp.29-42.
- 19- AL-Shimmery MN, AL-Khafaji AA and AL-Hilaly HA. (2011) Seroprevalence of CMV & *Toxo. gondii* in aborted women in AL-Diwaniyah province. *Q.M.J.*, **7**(11) pp.160-166.
- 20- Mumtaz C. Neval MD, Sirin MD, Nisel MD, Yeser K, Derici MD, Arzu MD. (2017) Seroprevalence of *T. gondii*, Rubella virus and CMV in pregnant women and the avidity assays importance. *Saudi. Med. J.*, **38** (7) pp.727-732.
- 21- Karabulut A, Türk M, Polat Y and Işık Balcı Y. (2011) Evaluation of Rubella, *T. gondii* and CMV seroprevalence between pregnant women in Denizli. *Turk. Med. Sci. J.*, **41** PP. 159-164.
- 22- Cim N, Erdin BN, Pariak M, Guven A and Bayram Y. (2015) seroprevalence of *T. gondii*, Rubella & CMV in pregnant women in Van. *Obset. Gynecol. Turk. J.*, **12**(2) PP. 79-82.
- 23- Naouri B, Bekhit R, Ahmed H, Mohsni E, Teleb N. (2011) Progress Toward Measles Elimination in the Eastern Mediterranean Region. *Infect. Dis. J.*, **204**(1 1) PP. 289-298.
- 24- Majeed HM, Alsamarai AM, Alsalihi FG, Alobaidi AH and Aljumaili ZKM. (2016) Seroprevalence of Rubella among Women with BOH. *Immuno. Virol. J.*, **1**(2) PP. 5-7.
- 25- Karad D and Kharat A . (2015) Seroprevalence of Torch in Bad Obstetrics History among HIV and Non-HIV Women in Solapur of Maharashtra\ India. *Hum. Virol. Retrovirol. J.*, **2** pp. 67.

## انتشار طفيلي المقوسة الكوندية و فيروس المضخم للخلايا وفيروس الحصبة الألمانية في النساء

### المجھضات في مدينة سامراء

سيناء ناجي محسن

كلية طب الاسنان ، جامعة تكريت ، تكريت ، العراق

#### الملخص

هدفت الدراسة الحالية تقييم نسبة الإصابة (مصلياً) لطفيلي المقوسات الكوندية والفيروس المضخم للخلايا وفيروس الحصبة الألمانية لـ 158 امرأة مجھضة في مستشفى سامراء العام خلال الفتره من كانون الثاني إلى كانون الأول 2016. حيث كانت نسبة الإصابة لكل من طفيلي المقوسة الكوندية 14,6%، وفيروس الحمى السوداء 10,8%، فيروس الحصبة الألمانية 8,2%، إصابة مشتركة لكل من طفيلي داء القطط وفيروس المضخم للخلايا 9,5%. فضلاً عن ذلك فقد بينت الدراسة الحالية إن الإصابة بطفيلي المقوسة الكوندية وفيروس الحمى السوداء وفيروس الحصبة الألمانية كانت الأعلى (24,2%، 15,2%، 15,2% على التوالي) في المجموعة العمرية 21-30 سنة، بينما الإصابة المشتركة لكل من طفيلي المقوسة الكوندية وفيروس الحمى السوداء كانت الأعلى (11,8%) في المجموعة العمرية 14-20 سنة. بينما كانت النسبة المؤية لعدد المقوسة الكوندية IgM 7%، IgG 3,2% و IgG & IgM 4,4%، والنسبة المؤية لعدد فيروس المضخم IgM 3,8%، IgG 1,9%، و IgG & IgM 2,5%. أما النسبة المصلية للإصابة المشتركة لكل من طفيلي المقوسة الكوندية وفيروس المضخم للخلايا فقد كانت ضد المقوسة الكوندية IgG وضد فيروس المضخم للخلايا IgM 0,63%، ضد المقوسة الكوندية IgM وضد المضخم للخلايا IgG 2,5%، ضد المقوسة الكوندية IgG وضد فيروس المضخم للخلايا IgM & IgG 1,9%، ضد المقوسة الكوندية IgM و ضد فيروس المضخم للخلايا IgG & IgM 3,2% و ضد المقوسة الكوندية IgM و ضد فيروس المضخم للخلايا IgG 1,3%.